## 1.2 | Constant and Linear Functions

Constant Functions: A constant function is of the form

$$f(x) = b$$

where b is a real number. The domain of a constant function is  $(-\infty, \infty)$ . A constant function will look like a horizontal line on the graph and intersect the y-axis at b.

**Linear Functions**: A linear function is of the form

$$f(x) = mx + b$$


where m and b are real numbers and  $m \neq 0$ . The domain of a linear function is  $(-\infty, \infty)$ .

**Intercepts**: The *y-intercept* of a function is where the function crosses the *y*-axis. This point will look like (0, f(0)). The *x-intercept* of a function is where the function crosses the *x*-axis. This point will look like (c, 0) for some real number c.

**Slope**: The slope of a linear function is the coefficient m. It is often described as *rise over run*. If you are given two points  $(x_1, y_1)$  and  $(x_2, y_2)$  from a linear function you can find the slope with the following formula:

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

1. Worked Example: Graph the function, find the slope and axis intercepts, if any: f(x) = 3x - 3




Scan the QR code for a video solution


## Problem Solving Tip 1. Finding axis intercepts:

To find an axis intercept, set the variable of the opposite axis equal to zero and solve. To find the y-axis intercept, set x = 0, and to find the x-axis intercept(s), set y = 0.

2. Graph the function, find the slope and axis intercepts, if any: f(x) = 2x - 1



3. Graph the function, find the slope and axis intercepts, if any: g(t) = 3 - t



Materials in PAL are not a suitable replacement for materials in class. These materials are not for use on exams.